

Rowan University

Rowan Digital Works

Cooper Medical School of Rowan University
Capstone Projects

Cooper Medical School of Rowan University

2020

Analysis of Risk Factors Predictive of Postoperative Complications in Hip Fracture Patients

Christopher Rivera-Pintado

Follow this and additional works at: https://rdw.rowan.edu/cmsru_capstones

Let us know how access to this document benefits you - share your thoughts on our feedback form.

Recommended Citation

Rivera-Pintado, Christopher, "Analysis of Risk Factors Predictive of Postoperative Complications in Hip Fracture Patients" (2020). *Cooper Medical School of Rowan University Capstone Projects*. 45.
https://rdw.rowan.edu/cmsru_capstones/45

This Poster is brought to you for free and open access by the Cooper Medical School of Rowan University at Rowan Digital Works. It has been accepted for inclusion in Cooper Medical School of Rowan University Capstone Projects by an authorized administrator of Rowan Digital Works. For more information, please contact brush@rowan.edu.

Introduction

Hip fracture management is an ongoing challenging task that orthopedic surgeons face daily. As the population is increasingly getting older, the incidence of this injury is becoming more prevalent with an estimation of 4.5 million by 2050 [1-2]. Patients who suffer hip fractures have a high one-year morbidity and mortality rate. Further, delay to surgery greater than 24 hours causes a further increase in these rates [3-8]. One of the factors for this delay is obtaining pre-operative cardiac evaluation and echocardiograms. In an attempt to prevent postoperative complications, these resources are often overused leading to an unnecessary delay to surgery [9-12]. This is a consequence of the unavailability of evidence about specific risk factors that are predictive of postoperative complications in hip fracture patients.

Objectives

- **Primary objective:** Evaluate how pre-operative cardiac evaluation affects time to surgery, in-hospital, and 30-day postoperative complications
- **Secondary objective:** Identify specific risk factors predictive of in-hospital, and 30-day postoperative complications

Methods

- Case-control study conducted in a teaching hospital located in Camden, New Jersey from January 2011 to December 2015
- Inclusion criteria: Age > 50 at the time of fracture, patients treated for a hip fracture
- Exclusion criteria: Age < 50 at the time of fracture, pathological fractures (i.e infection and cancer in the femur), pediatric and pregnant patients
- Electronic records were evaluated for demographic data, cardiology consultation, echocardiogram results and in-hospital and 30-day postoperative complications
- MACE complications included: atrial fibrillation, heart failure, chest pain, myocardial infarction, pulseless electrical activity (PEA) arrest, and deaths due to cardiovascular etiology.
- Statistical analysis: Univariate and multivariate regression

Results

Table 1. Demographics

Total number of patients	411
Females	312 (75.9%)
Males	99
Mean age	80.98 (65-109)
Average ASA	2.88 (1-4)
Average EF %	58.16 (15 - 80)
Time to surgery (days)	1.75 (0-11)
Length of stay (days)	5.45 (2-63)

Table 2. Mean time to surgery (days)

(-) Cardiology	(+) Cardiology	P-value
1.56	2.04	0.001
(-) Echocardiogram	(+) Echocardiogram	
1.55	2.44	<0.001

Table 3. In-hospital Postoperative Complications (P-values)

Risk Factor	Total CV	OR (CI)	MACE OR (CI)	Pulmonary OR (CI)	Neurological OR (CI)	Deaths #	OR (CI)
Age	0.1130	0.4775	0.7108	0.0684	0.4746	0.0098	1.242 (1.001-1.541)
Time to Surgery	0.0668	0.0855	0.2701	0.6794			
Fracture pattern							
Femoral neck	0.7062	0.6553	0.6255	1.1038	0.7793		
Intertrochanteric	0.2694	0.5369	0.2759	1.1013	0.7962		
Subtrochanteric	0.3874	0.1674	0.8733	0.8831	0.3488		
Basiscervical	0.2006	0.9861	0.9858	0.9896	0.9881		
Subcapital	0.9904	0.9838	0.9837	0.9929	0.9920		
Pertrochanteric	0.8597	0.4906	0.4393	0.3075	0.7328		
Type of surgery							
ORIF	0.2845	0.4047	0.4047	0.0302	0.443 (0.212-0.920)	0.2197	
Total Arthroplasty	0.5538	0.5538	0.9823	0.9868	0.3698		
Hemiarthroplasty	0.6678	0.4858	-	-	-		
Percutaneous Fixation	0.9875	0.9875	0.9875	0.9906	0.9853		
Intra-medullary Nail	0.1126	0.1999	0.9875	0.9906	0.9893		
ASA score	0.7902	0.1544	0.1591	0.3262	0.0160	3.585 (1.209-10.190)	
(+) CV consult	0.3063	0.0810	0.1161	0.5025	0.0093	2.500 (1.023-6.211)	
Risk Stratification							
Low	0.7825	0.9720	0.6874	0.5097	0.9761		
Moderate	0.6638	0.5839	0.0677	0.7751	0.3305		
High	0.9878	0.7523	0.0867	0.0010	5.038 (1.528-13.174)	0.9802	
(+) Echocardiogram	0.3407	0.2455	0.1502	0.1899	0.6880		
Ejection fraction < 50	0.9741	0.9655	0.2278	0.4354	0.9700		
Ejection fraction < 60	-	-	-	0.0463	5.000 (1.026-24.397)	-	
Ejection Fraction	0.3397	0.6288	0.0257	0.9403 (0.892-0.952)	0.0504	0.9870	

Table 4. 30-Days Postoperative Complications (P-values)

Risk Factor	Total CV	OR (CI)	MACE OR (CI)	Pulmonary OR (CI)	Neurological OR (CI)	Deaths #	OR (CI)
Age	0.1249	0.1249	0.5818	0.4102	0.2286		
Time to Surgery	0.5989	0.5989	0.4732	0.6716	0.2435		
Fracture pattern							
Femoral neck	0.8489	0.8489	0.8985	0.2910	0.9702		
Intertrochanteric	0.8424	0.8424	0.7194	0.9895	0.3325		
Subtrochanteric	0.9715	0.9715	0.9663	0.9807	0.9745		
Basiscervical	0.9912	0.9912	0.9895	0.0120	20.0 (1.08-381.803)	0.9881	
Subcapital	0.9961	0.9961	0.9953	0.9957	0.9947		
Pertrochanteric	0.9794	0.9794	0.0385	4.08 (1.12-15.84)	0.9774	0.9723	
Type of surgery							
ORIF	0.2302	0.2302	0.4864	0.3515	0.7910		
Total Arthroplasty	0.9888	0.9888	0.9867	0.0365	11.3 (1.16-109.886)	0.9853	
Hemiarthroplasty	-	-	-	-	-		
Percutaneous Fixation	0.9921	0.9921	0.9906	0.9913	0.9894		
Intra-medullary Nail	0.9921	0.9921	0.9906	0.9913	0.9894		
ASA score	0.8472	0.8472	0.8083	0.8382	0.7470		
(+) CV consult	0.9875	0.9875	0.9819	0.9946	0.9528		
Risk Stratification							
Low	0.9823	0.9823	0.9790	0.7109	0.9762		
Moderate	0.9833	0.9833	0.1511	0.9598	0.9872		
High	0.0077	12.2 (2.1-77.1)	0.0077	12.2 (2.1-77.1)	0.9775	0.9752	0.9745
(+) Echocardiogram	0.8878	0.8878	0.2134	0.5327	0.9003		
Ejection fraction < 50	0.9742	0.9742	0.9700	0.9742	0.9742		
Ejection Fraction	0.9416	0.9416	0.5604	0.6534	0.9416		

Conclusion

- Obtaining cardiacology consultation and echocardiogram alone delay surgery by 0.48 and 0.89 days, respectively
- In the context of cardiac risk stratification, high risk stratification was a predictor for 30-day postoperative cardiac complications and in-hospital PEA arrest
- Cardiac risk stratification and echocardiogram ejection fraction were not predictive of any other postoperative cardiac complications
- Obtaining a cardiac evaluation and high ASA scores, independently, resulted as predictor of in-hospital deaths
- Delay to surgery was predictive for in-hospital PEA arrest and deaths
- Pertrochanteric fractures were predictive of 30-day postoperative pulmonary complications.

Discussion

Our data demonstrates important predictors of in-hospital and 30-day complications. Effort should be made to minimally delay time to surgery in patients with hip fractures. The risk factors analyzed in this study should be taken into consideration during the decision making process in order to analyze the benefits and risk association with early surgical fixation. We encourage researchers to continue analyzing risk factors predictive of postoperative complications to further improve the decision making process and prevent the overuse of resources during the process.

References

1. Brauer CA, Coca-Perrallon M, Cutler DM, et al. Incidence and mortality of hip fractures in the United States. JAMA. 2009;302:1573-1579.
2. Gulberg B, Johnell O, Kanis JA. World-wide projections for hip fracture. Osteoporos Int 1997;7(5):407-13
3. Moran CG, Wenn RT, Siskand M, Taylor AM (2005) Early mortality after hip fracture: is delay before surgery important? J Bone Joint Surg Am 87(10):483-489
4. Hamlet WPF, Lieberman JR, Freedman EL, Dorey FJ, Fletcher A, Johnson EE (1997) Influence of health status and the timing of surgery on mortality in hip fracture patients. Am J Orthop 26:621-627
5. Villar RM, Allen SM, Barnes SJ (1986) Hip fractures in healthy patients: operative delay versus prognosis. Br Med J (Clin Res Ed) 293:1203-1204
6. Bredahl C, Nyholm B, Hindsholm FB, Mortensen JS, Olesen AS (1992) Mortality after hip fracture: results of operation within 12 h of admission. Injury 23:83-86
7. Zuckerman JD, Skovron ML, Kovalev KI, Aharonoff G, Frankel VH (1995) Postoperative complications and mortality associated with operative delay in older patients who have a fracture of the hip. J Bone Joint Surg Am 77:1551-1556
8. Kamezakar H, Ayhan E, Urali MC, Seker A, Karaca S (2010) Predictors of mortality in elderly patients with an inter-trochanteric or a femoral neck fracture. J Trauma 68(1):153-158
9. Orszag GM, Hannan EL, Magaziner J, et al. Hip fracture in the older patient: reasons for delay in hospitalization and timing of surgical repair. J Am Geriatr Soc 2002;50:1338-1340
10. Siligen A, Poludniak K, Dulaney-Cripe E, et al. Adherence to preoperative cardiac clearance guidelines in hip fracture patients. J Orthop Trauma. 2015;29:500-503.
11. Abbas K, Umer M, Akbari R. Preoperative cardiac evaluation in proximal femur fractures and its effects on the surgical outcome. Acta Orthop Traumatol Turc. 2012;46(4):250-4.
12. Luttrell K, Nana A. Effect of preoperative transthoracic echocardiogram on mortality and surgical timing in elderly adults with hip fracture. J Am Geriatr Soc. 2015 Dec;63(12):2605-9.