

Rowan University

Rowan Digital Works

Rowan-Virtua Research Day

25th Annual Research Day

May 6th, 12:00 AM

A Peculiar Presentation of Cardiac Sarcoidosis As Third-Degree Atrioventricular Heart Block Complicated By Right Ventricle Perforation

Yash Suthar
Rowan University

Pooja Patel
Rowan University

Nikhita Kothari
Rowan University

Kristofer Hillegas
Virtua Cardiology

Follow this and additional works at: https://rdw.rowan.edu/stratford_research_day



Part of the [Cardiovascular Diseases Commons](#), and the [Pathological Conditions, Signs and Symptoms Commons](#)

Let us know how access to this document benefits you - share your thoughts on our [feedback form](#).

Suthar, Yash; Patel, Pooja; Kothari, Nikhita; and Hillegas, Kristofer, "A Peculiar Presentation of Cardiac Sarcoidosis As Third-Degree Atrioventricular Heart Block Complicated By Right Ventricle Perforation" (2021). *Rowan-Virtua Research Day*. 2.
https://rdw.rowan.edu/stratford_research_day/2021/may6/2

This Poster is brought to you for free and open access by the Conferences, Events, and Symposia at Rowan Digital Works. It has been accepted for inclusion in Rowan-Virtua Research Day by an authorized administrator of Rowan Digital Works.

A Peculiar Presentation of Cardiac Sarcoidosis As Third-Degree Atrioventricular Heart Block Complicated By Right Ventricle Perforation

ROWAN UNIVERSITY



Yash Suthar, DO¹; Pooja Patel, OMS-III²; Nikhita Kothari, DO³; Kristofer Hillegas, DO, FACC⁴

Dept. of Cardiovascular Medicine¹, Rowan University – School of Osteopathic Medicine²; Dept. of Internal Medicine³; Virtua Cardiology - Central⁴

BACKGROUND

- Cardiac sarcoidosis is a rare, inflammatory multisystem disorder that manifests as noncaseating granulomas of multiple organs.
- The clinical presentation of sarcoidosis is variable and may be underrecognized by clinicians.
- Individuals may be asymptomatic during their lifetimes and may be incidentally diagnosed with the disease for the first-time during autopsy.
- Patients with cardiac involvement of sarcoidosis may initially present with complications including arrhythmias, heart block, bundle branch block, congestive heart failure, pericardial effusion, pulmonary hypertension, and/or sudden cardiac death.

CASE DESCRIPTION

- A 59-year-old African American female with past medical history of untreated hypertension and hyperlipidemia presented following multiple episodes of syncope and near syncope.
- On initial exam, patient was found to be in 3rd degree atrioventricular heart block with a ventricular escape rhythm of 40 beats per minute. **(Figure 1)**
- A temporary transvenous pacemaker was placed through the right Internal Jugular Vein with good capture.
- The patient was transferred to a tertiary care center for a permanent pacemaker placement.
- Upon arrival, the patient was noted to have regained intrinsic rhythm with EKG demonstrating normal sinus rhythm with a first-degree atrioventricular heart block and right bundle branch block at 70 bpm. **(Figure 2)**
- Later that day, the temporary pacemaker malfunctioned and stopped capturing despite increasing the pacing threshold to the maximum setting.
- CT scan of the chest demonstrated that patient's transvenous pacemaker lead had perforated through the free wall of the right ventricle with the tip of the pacing catheter extending approximately 3-4 cm into the pericardium. **(Figures 5 & 6)**
- The original temporary pacemaker was then removed under transesophageal echocardiogram guidance and a separate temporary pacemaker was floated through the right femoral vein under fluoroscopic guidance.
- A chest x-ray performed initially showed bilateral hilar adenopathy and the CT scan of the chest verified the presence of mediastinal, subcarinal and hilar adenopathy concerning for pulmonary sarcoidosis. **(Figures 3 & 4)**




Figure 1
EKG demonstrating 3rd degree atrioventricular heart block with a ventricular escape rhythm of 40 bpm.




Figure 2
EKG demonstrating normal sinus rhythm with first degree atrioventricular heart block and right bundle branch block.




Figure 3
CXR showing bilateral peri-hilar lymphadenopathy.




Figure 4
Chest CT scan showing bilateral peri-hilar lymphadenopathy.




Figure 5
Chest CT showing perforation of right ventricle by temporary transvenous pacemaker lead.




Figure 6
Chest CT coronal slice showing perforation of right ventricle by temporary transvenous pacemaker lead.

CASE OUTCOME

- A dual chamber implantable cardioverter defibrillator was placed in accordance with the 2017 American College of Cardiology/ American Heart Association/ Heart Rhythm Society (AHA/ACC/HRS) guidelines.
- Patient was discharged in a stable condition to follow up with a cardiologist and a pulmonologist for further confirmatory testing.

DISCUSSION

- The patient's clinical presentation with episodes of heart block and pulmonary lymphadenopathy lead to the presumptive diagnosis of cardiac sarcoidosis.**
- Cardiac Sarcoidosis is associated with a poor prognosis due to the many lethal complications arising from the disease, particularly high degree atrioventricular blocks, ventricular arrhythmias, and sudden cardiac death.
- A diagnosis of cardiac sarcoidosis can be made by non-invasive testing such as a cardiac MRI or a cardiac FDG-PET scan if a patient has histologic or clinical evidence of extracardiac sarcoidosis and signs and symptoms of palpitations, syncope/near syncope, conduction abnormalities such as heart block, ventricular tachycardia, or even echocardiographic abnormalities such as regional wall motion abnormality, ventricular aneurysm, basal septal thinning, or depressed LV ejection fraction.
- It is recommended that patients with extracardiac sarcoidosis with or without cardiac symptoms should be evaluated for subclinical or clinical cardiac involvement.

CONCLUSION

- This case highlights a severe, insidious clinical manifestation of cardiac sarcoidosis and the importance of early recognition and treatment of the condition.

REFERENCES

- Fleming, H., *Sarcoid heart disease*. British Heart Journal, 1974. **36**: p. 54-68
- Baughman RP, T.A., Judson MA, Rossman MD, et al. , *Clinical characteristics of patients in a case control study of sarcoidosis*. . Am J Respir Crit Care Med., 2001. **164**: p. 1885–1889.
- Mehdi, M.R., F. Machado. Dean, Schraufnagel. Nadera, J. Sweiss. Robert P. Baughman and M.M. Mirsaedi, Roberto F. Schraufnagel, Dean. Sweiss, Nadera. Baughman, Robert., *Racial Differences in Sarcoidosis Mortality in the United States*. Chest, 2015. **147**(2): p. 438.
- Yazaki Y, I.M., Hiroe M, et al., *American Journal of Cardiology*. 2001, Central Japan Heart Study Group Prognostic determinants of long-term survival in Japanese patients with cardiac sarcoidosis treated with prednisone. . **88**(9): p. 1006-1010.