

Rowan University

Rowan Digital Works

Rowan-Virtua Research Day

26th Annual Research Day

May 5th, 12:00 AM

Factors Associated with Unplanned Conversion to Open in Nephrectomy for Kidney Cancer

Young Son

Rowan University

Benjamin A. Fink

Rowan University

Justine Garfinkel

Rowan University

Lance Earnshaw

Rowan University

Brian Thomas

Rowan University

See next page for additional authors

Follow this and additional works at: https://rdw.rowan.edu/stratford_research_day



Part of the [Nephrology Commons](#), [Surgery Commons](#), [Surgical Procedures](#), [Operative Commons](#), and the [Urogenital System Commons](#)

Let us know how access to this document benefits you - share your thoughts on our [feedback form](#).

Son, Young; Fink, Benjamin A.; Garfinkel, Justine; Earnshaw, Lance; Thomas, Brian; Mueller, Thomas; and Sussman, David, "Factors Associated with Unplanned Conversion to Open in Nephrectomy for Kidney Cancer" (2022). *Rowan-Virtua Research Day*. 65.

https://rdw.rowan.edu/stratford_research_day/2022/May5/65

This Poster is brought to you for free and open access by the Conferences, Events, and Symposia at Rowan Digital Works. It has been accepted for inclusion in Rowan-Virtua Research Day by an authorized administrator of Rowan Digital Works.

Submitting Author(s)

Young Son, Benjamin A. Fink, Justine Garfinkel, Lance Earnshaw, Brian Thomas, Thomas Mueller, and David Sussman

Factors Associated with Unplanned Conversion to Open in Nephrectomy for Kidney Cancer

Young Son, D.O., Benjamin A. Fink OMS3, Justine Garfinkel OMS3, Lance Earnshaw OMS4, Brain Thomas OMS3, Thomas Mueller, M.D., David Sussman, D.O.

Jefferson New Jersey Urology, 18 E. Laurel Rd., Stratford, NJ 08084

Background

Minimally invasive surgery (MIS) has been adopted as an approach in kidney surgery. Laparoscopic kidney surgery has been introduced in the 1990s with robotics emerging a decade after.^{1,2} The minimally invasive approach has been technically feasible and has been shown to be noninferior with preserved oncology standards to open surgery.³ The ubiquitous use of MIS for kidney cancer has been standard of practice; however, unplanned conversion to open kidney surgery has been characterized at 4.9% for laparoscopic radical nephrectomy compared to 6.0% in robotic radical nephrectomy.⁴ Another analysis of 54,246 patients undergoing partial nephrectomy for kidney cancer observed an unplanned open conversion rate of 2.87% for cT1 renal masses. Furthermore, the unplanned conversion to open radical or partial nephrectomy after an attempted minimally invasive approach has been an independent predictor of increased risk of 30-day hospital readmission.⁵ Currently, the relative risk factors to predict unplanned conversion to open surgery has not been well characterized. Greater understanding of risk factors for unplanned open conversion has clinical implications to reduce intraoperative and postoperative outcomes. The early recognition of nationwide risk and predictors may aid in identifying patients for planned open kidney surgery. We aim to use the American College of Surgeons National Surgical Quality Improvement Program (ACS-NSQIP) database to identify predictors and outcomes in a contemporary cohort.

Methods

The 2019 and 2020 NSQIP data was analyzed in a retrospective cohort study of radical cystectomy patients. A total of 14,186 patients were evaluated and 4,862 patients met the inclusion criteria of nephrectomy performed for kidney cancer. The groups were then subcategorized to planned MIS only (endoscopic, laparoscopic, other MIS, and robotic) (n=4756) and unplanned conversion to open starting as either robotic, laparoscopic, or endoscopic (n=106). Wilcoxon signed-rank test was utilized for continuous variables as the data was not normally distributed. For categorical variables, Fisher's exact test was performed and Chi-squared analysis where more than two categories were evaluated. Statistical significance was set at P-value < 0.05.




Figure 1. CONSORT flow diagram illustrates the composition of the cohort.

Tables

Table 1. Patient Characteristics, Unplanned Conversion to Open Nephrectomy

	Total Cohort n = 4862	Unplanned Conversion to Open n = 106	Minimally Invasive Surgery n = 4756	P Value
Patient Demographics				
Mean Age (Range)	61.28 (19 - 89)	62.58 (32 - 88)	61.25 (19 - 89)	
Male Gender (%)	3170 (65.20%)	72 (67.92%)	3098 (65.14%)	
Non-Caucasian Race (%)	1580 (32.50%)	39 (36.79%)	1541 (32.40%)	
Hispanic Ethnicity (%)	353 (7.26%)	4 (3.77%)	349 (7.34%)	
Preoperative Considerations:				
Admitted from Other Than Home (%)	23 (0.4%)	2 (1.9%)	21 (0.4%)	
Preoperative ASA Classification Greater than 3 (%)	3008 (61.9%)	80 (75.4%)	2928 (61.6%)	P < 0.01
Preoperative Non-Functional Health Status (%)	40 (0.82%)	0 (0.0%)	40 (0.8%)	
>10% Decrease in Body Weight in 6 Months (%)	66 (1.4%)	4 (3.8%)	62 (1.3%)	P = 0.02
Preoperative Mechanical Bowel Prep (%)	754 (15.5%)	11 (10.4%)	743 (15.6%)	
Preoperative Oral Antibiotic Prep (%)	70 (1.4%)	1 (0.9%)	69 (1.5%)	
Prior Pelvic Surgery (%)	1608 (33.1%)	41 (38.7%)	1567 (32.9%)	
Prior Pelvic Radiotherapy (%)	75 (1.5%)	0 (0.0%)	75 (1.6%)	
Chemotherapy (within 90 days of surgery) (%)	103 (2.1%)	3 (2.8%)	100 (2.1%)	
ASA Classification:				
Class 1 (%)	62 (1.3%)	1 (0.9%)	61 (1.3%)	P < 0.01
Class 2 (%)	1792 (36.9%)	25 (23.6%)	1767 (37.2%)	
Class 3 (%)	2834 (58.3%)	71 (67.0%)	2763 (58.1%)	
Class 4 (%)	171 (3.5%)	8 (7.5%)	163 (3.4%)	P = 0.02
TNM Classification:				
T1, T1a, T1b (%)	3481 (71.6%)	57 (53.8%)	3424 (72.0%)	P < 0.01
T2, T2a, T2b (%)	334 (6.9%)	8 (7.5%)	326 (6.8%)	
T3, T3a, T3b, T3c (%)	1042 (21.4%)	32 (30.2%)	1010 (21.2%)	P = 0.03
T4 (%)	33 (0.7%)	9 (8.5%)	24 (0.5%)	P < 0.01
Nx (%)	4199 (86.4%)	74 (69.8%)	4125 (86.7%)	P < 0.01
N0 (%)	472 (9.7%)	21 (19.8%)	451 (9.5%)	P < 0.01
N1 (%)	61 (1.3%)	4 (3.8%)	57 (1.2%)	P = 0.02
N2 (%)	2196 (45.2%)	44 (41.5%)	2152 (45.2%)	
N3 (%)	67 (1.4%)	5 (4.7%)	62 (1.3%)	P < 0.01
Pre-Operative Lab Values:				
Elevated Creatinine (Greater than 1.6 mg/dL) (%)	337 (6.9%)	15 (14.2%)	322 (6.8%)	P < 0.01
Elevated BUN (Greater than 21 mg/dL) (%)	1155 (23.8%)	33 (31.1%)	1122 (23.6%)	
Elevated PTT (Greater than 36 seconds) (%)	233 (4.8%)	13 (12.3%)	220 (4.6%)	P < 0.01
Elevated INR (Greater than 1.6) (%)	271 (5.6%)	12 (11.3%)	259 (5.5%)	P < 0.01
Decreased Platelet Count (Less than 149 x 10 ⁹ /Liter) (%)	567 (11.7%)	17 (16.0%)	550 (11.6%)	
Comorbidities:				
Transfusion within 72 hours of Surgery (%)	15 (0.3%)	2 (1.9%)	13 (0.3%)	P < 0.01
Currently on Hemodialysis (%)	124 (2.6%)	8 (7.5%)	116 (2.4%)	P < 0.01
Current Smoker (within past year) (%)	789 (16.2%)	13 (12.3%)	776 (16.3%)	
Diabetes Mellitus Treated with Insulin (%)	342 (7.0%)	8 (7.5%)	334 (7.0%)	
Diabetes Mellitus Treated with Oral Medication (%)	754 (15.5%)	18 (17%)	736 (15.5%)	
Disseminated Cancer (%)	155 (3.2%)	10 (9.4%)	145 (3.0%)	P < 0.01
History of Severe COPD (%)	184 (3.8%)	2 (1.9%)	182 (3.8%)	
Hypertension (Treated with Medication) (%)	2984 (61.4%)	69 (65.1%)	2915 (61.3%)	
Steroid Use for Chronic Medical Condition (%)	195 (4.0%)	5 (4.7%)	190 (4.0%)	
Dyspnea (%)	285 (5.9%)	12 (11.3%)	273 (5.8%)	P = 0.03
Bleeding Disorder (%)	124 (2.6%)	6 (5.7%)	118 (2.5%)	P = 0.04

P Value represents T-test for continuous variables or Chi-Square for categorical variables.
ASA = American Society of Anesthesiologists, COPD = Chronic Obstructive Pulmonary Disorder, BUN: Blood urea nitrogen

Table 2. Postoperative Outcomes Frequency Table, Unplanned conversion to open

Outcome Variables	Total Cohort n = 4862	Unplanned Conversion to Open n = 106	Minimally Invasive Surgery n = 4756	P Value
Readmission to Hospital (%)	227 (4.7%)	17 (16.0%)	210 (4.4%)	P < 0.01
Pt Requiring Bleeding Transfusion (%)	189 (3.9%)	35 (33.0%)	154 (3.2%)	P < 0.01
Mean Total Operating Time in Minutes (Range)	176.4 (4 - 769)	255.08 (66 - 732)	174.7 (4 - 769)	P < 0.01
Mean Length of Hospital Stay (Range)	2.4 (0 - 36)	5.07 (1 - 27)	2.3 (0 - 36)	P < 0.01
Mean Days from Operation to Discharge (Range)	2.1 (0 - 30)	0.85 (0 - 21)	2.2 (0 - 30)	P < 0.01
Superficial Incisional Surgical Site Infection (%)	62 (1.3%)	3 (2.8%)	59 (1.2%)	
Deep Incisional Surgical Site Infection (%)	7 (0.1%)	2 (1.9%)	5 (0.1%)	P < 0.01
Organ Space Incisional Surgical Site Infection (%)	27 (0.6%)	6 (5.7%)	21 (0.4%)	P < 0.01
Sepsis (%)	20 (0.4%)	2 (1.9%)	18 (0.4%)	P = 0.02
Septic Shock (%)	10 (0.2%)	1 (0.9%)	9 (0.2%)	
Urinary Tract Infection (%)	52 (1.1%)	0 (0.0%)	52 (1.1%)	
Acute Renal Failure (%)	15 (0.3%)	4 (3.8%)	11 (0.2%)	P < 0.01
Pneumonia (%)	54 (1.1%)	1 (0.9%)	53 (1.1%)	
CVA/Stroke with Neurological Deficit (%)	6 (0.1%)	0 (0.0%)	6 (0.1%)	
Cardiac Arrest Requiring CPR (%)	16 (0.3%)	1 (0.9%)	15 (0.3%)	
Myocardial Infarction (%)	26 (0.5%)	1 (0.9%)	25 (0.5%)	
Pulmonary Embolism (%)	24 (0.5%)	3 (2.8%)	21 (0.4%)	P < 0.01
DVT/Thrombophlebitis (%)	26 (0.5%)	4 (3.8%)	22 (0.5%)	P < 0.01
Discharge to SNF (%)	119 (2.5%)	11 (10.4%)	108 (2.3%)	
Discharge to Home (%)	4743 (97.6%)	95 (89.6%)	4648 (97.7%)	
Rectal Injury (%)	3 (0.1%)	0 (0.0%)	3 (0.1%)	
Unplanned Intubation (%)	18 (0.4%)	0 (0.0%)	18 (0.4%)	
Wound Disruption (%)	17 (0.4%)	4 (3.8%)	13 (0.3%)	P < 0.01
On Ventilator Greater than 48 Hours (%)	17 (0.3%)	2 (1.9%)	15 (0.3%)	P < 0.01
Clostridium Difficile Colitis (%)	10 (0.2%)	1 (0.9%)	9 (0.2%)	
Prolonged NPO or NGT Use (%)	82 (1.7%)	13 (12.3%)	69 (1.5%)	P < 0.01
Lymphocele/Lymphatic Leak (%)	82 (1.7%)	6 (5.7%)	76 (1.6%)	P < 0.01
Unplanned Reoperation (%)	85 (1.7%)	10 (10.9%)	75 (1.6%)	P < 0.01

P Value represents T-Test for continuous variables and Chi-Square for categorical variables.
UTI = Urinary Tract Infection, PNA = Pneumonia, DVT = Deep Vein Thrombosis, SNF = Skilled Nursing Facility, NPO = Nothing by mouth, NGT = Nasogastric Tube

Results

The unplanned conversion to open rate was 2.2%. The age (mean (SD)) of the cohort was 62.6 (11.2) and 61.3 (12.2) for the unplanned and the planned group. There was a baseline difference with preoperative labs including BUN, creatinine, PTT, and INR. An increased incidence of T3 and T4 (30.2% and 8.5%) in the unplanned group compared to the (21.2% and 0.50%) in the planned group was noted. ASA classification 4 also favored the unplanned group vs. planned group (Table 1). The outcome variables were also compared between the unplanned and the planned group (Table 2).

Discussion

Based on our study, multiple factors were associated with conversion to open in nephrectomy. Higher tumor stage, ASA classification, kidney function, and PT/INR were all predictive factors of conversion to open in nephrectomies. The open group had worse clinical outcomes including length of hospital stay, discharge to nursing care, prolonged NPO/NGT, and lymph leak.

Limitations

This study was performed retrospectively and, therefore, no causality can be drawn from these findings. Another limitation is surgeon preference when converting to open nephrectomy. In these cases, each urological surgeon may have a different threshold for open conversion, and this variable cannot be controlled for. Although we performed a multivariable statistical regression, unaccounted for confounding variables, other than primary surgeon, could potentially influence the data.

References

- Miller DC, Saigal CS, Banerjee M, Hanley J, Litwin MS, Urologic Diseases in America Project. Diffusion of surgical innovation among patients with kidney cancer. *Cancer* 2008;112:1708-1717.
- Gill IS, Sung GT, Hsu TH, Meraney AM. Robotic remote laparoscopic nephrectomy and adrenalectomy: The initial experience. *J Urol* 2000;164:2082-2085.
- Golombos DM, Chughtai B, Trinh QD, Thomas D, Mao J, Te A, O'Malley P, Scherr DS, Del Pizzo J, Hu JC, Sedrakyan A. Minimally invasive vs open nephrectomy in the modern era: does approach matter? *World J Urol*. 2017 Oct;35(10):1557-1568. doi: 10.1007/s00345-017-2040-6. Epub 2017 May 5. PMID: 28477204
- Khanna A, Campbell SC, Murthy PB, Ericson KJ, Nyame YA, Abouassaly R. Unplanned Conversion from Minimally Invasive to Open Kidney Surgery: The Impact of Robotics. *J Endourol*. 2020 Sep;34(9):955-963. doi: 10.1089/end.2020.0357. Epub 2020 Aug 6. PMID: 32597204.
- Razdan S, Okhawere K, Wilson M, Nkemdirim W, Korn T, Meilika K, Badani K. Conversion to Open Radical or Partial Nephrectomy Associated with Unplanned Hospital Readmission After Attempted Minimally Invasive Approach. *J Laparoendosc Adv Surg Tech A*. 2021 Dec 28. doi: 10.1089/lap.2021.0537. Epub ahead of print. PMID: 34962141.