

7-2011

Modifiable Disease Risk, Readiness to Change, and Psychosocial Functioning Improve With Integrative Medicine Immersion Model

Ruth Q. Wolever

Daniel M. Webber

Justin P. Meunier

Jeffrey M. Greeson

Rowan University, greeson@rowan.edu

Evangeline R. Lausier

See next page for additional authors

Follow this and additional works at: http://rdw.rowan.edu/csm_facpub



Part of the [Health Psychology Commons](#)

Recommended Citation

Wolever, Ruth Q.; Webber, Daniel M.; Meunier, Justin P.; Greeson, Jeffrey M.; Lausier, Evangeline R.; and Gaudet, Tracy W., "Modifiable Disease Risk, Readiness to Change, and Psychosocial Functioning Improve With Integrative Medicine Immersion Model" (2011). *Faculty Scholarship for the College of Science & Mathematics*. 75.
http://rdw.rowan.edu/csm_facpub/75

This Article is brought to you for free and open access by the College of Science & Mathematics at Rowan Digital Works. It has been accepted for inclusion in Faculty Scholarship for the College of Science & Mathematics by an authorized administrator of Rowan Digital Works. For more information, please contact jiras@rowan.edu, rdw@rowan.edu.

Authors

Ruth Q. Wolever, Daniel M. Webber, Justin P. Meunier, Jeffrey M. Greeson, Evangeline R. Lausier, and Tracy W. Gaudet



Published in final edited form as:

Altern Ther Health Med. 2011 ; 17(4): 38–47.

Modifiable Disease Risk, Readiness to Change, and Psychosocial Functioning Improve With Integrative Medicine Immersion Model

Ruth Q. Wolever, PhD, Daniel M. Webber, MS, Justin P. Meunier, BA, Jeffrey M. Greeson, PhD, Evangeline R. Lausier, MD, and Tracy W. Gaudet, MD

Ruth Q. Wolever, PhD, is director of research and health psychology at Duke Integrative Medicine, Duke University Medical Center, Durham, North Carolina, and assistant professor in the Department of Psychiatry and Behavioral Sciences, Duke University Medical Center. Daniel M. Webber, MS, was a clinical research coordinator at Duke Integrative Medicine and in the Department of Psychiatry and Behavioral Sciences, Duke University Medical Center. Justin P. Meunier, BA, was a research assistant at Duke Integrative Medicine. Jeffrey M. Greeson, PhD, is a health psychologist at Duke Integrative Medicine and an assistant professor in the Department of Psychiatry and Behavioral Sciences, Duke University Medical Center. Evangeline R. Lausier, MD, is the clinical director at Duke Integrative Medicine and an assistant professor in the Department of Medicine, Duke University Medical Center. Tracy W. Gaudet, MD, was the executive director at Duke Integrative Medicine and assistant professor in the Department of Obstetrics and Gynecology, Duke University Medical Center.

Abstract

Background—Stroke, diabetes, and coronary heart disease (CHD) remain leading causes of death in the United States and are largely attributable to lifestyle behaviors. Integrative medicine can provide a supportive partnership that focuses on improving health by identifying and implementing lifestyle changes based upon personal values and goals.

Objective—This prospective observational study was designed to assess the effectiveness of an integrative medicine intervention on modifiable disease risk, patient activation, and psychosocial risk factors for stroke, diabetes, and CHD.

Design—Sixty-three adults participated in a 3-day comprehensive, multimodal health immersion program at Duke Integrative Medicine, Duke University Medical Center, Durham, North Carolina. Participants received follow-up education, physician support, and telephonic health coaching between the immersion program and the endpoint 7 to 9 months later.

Primary Outcome Measures—Psychosocial functioning, readiness to change health behaviors, and risk of developing diabetes, stroke, and CHD were assessed at baseline and endpoint.

Results—Although cardiac risk remained unchanged ($P = .19$) during the study period, risk of diabetes ($P = .02$) and stroke ($P < .01$) decreased significantly. Perceived stress remained unchanged, but improvements were seen in mood ($P < .05$) and relationship satisfaction ($P < .004$). Patients became more activated towards self-management of health ($P < .001$), endorsed greater readiness to change health behaviors ($P < .01$), and reported increased aerobic exercise ($P < .001$) and stretching ($P = .006$) following the intervention.

Conclusion—An integrative health model can help patients become more engaged in self-management of health and support them in making and maintaining healthy lifestyle changes. These findings provide support for use of an integrative health model in adult disease risk reduction.

Stroke, diabetes, and coronary heart disease (CHD) remain leading causes of death in the United States¹ and are largely attributable to behavior.²⁻⁴ Expert recommendations for disease risk reduction include increasing physical activity, moderating caloric intake, and reducing stress.⁵⁻⁸ Nonetheless, inactivity remains pervasive in the United States, with as much as 62% of the population not participating in any vigorous activity in the span of a year.⁹ Caloric intake has increased steadily over the past 3 decades,¹⁰ and the resulting imbalance between caloric intake and energy expenditure has resulted in an increased rate of obesity and excess weight.¹ In addition, a large segment of the population reports negative psychosocial factors—chronic stress, negative mood, and low levels of social support—that contribute to risk of stroke, diabetes, and CHD.¹¹⁻¹⁴

Prospective trials have clearly demonstrated that improvements in diet¹⁵ and exercise¹⁶ can have a profound impact upon markers for disease risk. Furthermore, recent studies have shown that alternative approaches to stress reduction such as yoga^{17,18} and meditation¹⁹⁻²¹ can likewise influence chronic disease risk. Research has shown that improving health behaviors can lead to improvement in overall health and well-being; however, individuals desiring to better their health often face obstacles when it comes to initiating and maintaining changes in behavior. An integrative health model may help overcome these hurdles through supportive patient partnerships that focus on identifying and implementing lifestyle changes based upon personal values and goals. Integrative health professionals trained in coaching can support and promote this process by building trusting relationships with patients that encourage personal growth, enhance motivation, and promote self-efficacy.²²

There has been only limited investigation into the effectiveness of a patient-centered program that combines multiple strategies into a whole-person paradigm for disease risk reduction.^{23,24} Accordingly, the aim of the current study was to evaluate patient outcomes from an integrative health program designed to provide holistic, patient-centered care and incorporate conventional and complementary medicine approaches for the reduction of chronic disease risk.

METHODS

Study Design

A prospective observational study design was used to assess changes in modifiable disease risk, patient activation, and behavioral and psychosocial measures after an integrative medicine health program. This program included two principal components: a 3-day health immersion phase conducted at an academic integrative medicine facility and a support phase consisting of telephonic health coaching and monthly didactic sessions on health topics. The research protocol was approved by the Duke University Medical Center Institutional Review Board.

Participants

Participants included 63 individuals (33 male and 30 female) who provided written informed consent and attended a discounted, self-paid 3-day health immersion program at Duke Integrative Medicine. Participation was limited to members and employees of an early retirement community in South Carolina, all literate in English and 18 years of age or older. Individuals were excluded if they were pregnant, had severe psychiatric disease, a cognitive

impairment (eg, dementia, Alzheimer's disease), or other conditions that would limit their ability to provide informed consent.

Intervention

Phase I: 3-day Health Immersion—During the health immersion, participants spent 3 days at the Integrative Medicine center, participating in a variety of conventional and complementary therapies including nutritional counseling, acupuncture, massage, exercise training, yoga, and mind-body therapy (Table 1). Classes and individual consultations focused on topics identified on the Duke Integrative Medicine (IM) Wheel of Health (Figure 1).²⁵ This model centers on mindfulness and includes topics related to both professional care (eg, pharmaceuticals/supplements, preventive medicine, and conventional/CAM treatments) and self-care (eg, mind-body connection, movement/exercise, nutrition, physical environment, relationships, and personal growth/spirituality). A team of health care providers worked with each participant individually to develop a personalized health plan²⁶ corresponding to the 10 integral areas of health identified on the Wheel of Health (Figure 1). During the 3 days, participants also met with health coaches to reflect on personal goals and values as well as to refine and implement the health plan. Thus by the end of the immersion, participants had established a comprehensive multimodal plan for implementing lifestyle changes.

Phase II: Support Phase—After completing the health immersion, participants were provided with follow-up integrative health coaching (IHC) by telephone. IHC is a relatively recent addition to the field of coaching.²² The IHC model builds upon elements of life coaching such as appreciative inquiry²⁷ and motivational interviewing²⁸ and in addition helps patients explore motivation to move toward whole-person health as conceptualized by the Wheel of Health (Figure 1). Specifically, IHC strives to link desired lifestyle changes with personal values and mission.

Masters-level professionals trained in IHC performed the health coaching and provided participants with support and guidance in implementing their personalized health plans. The support phase included five 30- to 40-minute one-on-one phone calls during the 31- to 37-week period. Participants initiated calls at agreed-upon times convenient to both coaches and participants. Within the first 2 months of the support phase, participants also were offered a 30-minute phone consultation with their integrative physicians. This phone call provided an opportunity for physicians to address patient questions, update participants on baseline lab results, and explain results from Know Your Number (KYN) disease risk assessments (Biosignia, Inc, Durham, North Carolina; see Measures). Participants also were provided with eight monthly education lectures that were delivered in person by Duke IM clinicians in the participants' home communities. Lectures provided general preventive health education and further expanded upon the Wheel of Health concepts learned in the 3-day health immersion program (Table 2).

Measures

Disease Risk—Primary outcomes included 5-year risk of developing diabetes, stroke, and CHD, as estimated by the KYN disease assessment tool. The KYN tool calculates disease risk using a multivariate, meta-analytic disease model that incorporates biological measures and information provided by a comprehensive patient questionnaire (demographics, disease history, family history, and behavioral measures; Table 3).^{29,30} The model provides estimates of relative risk (5-year absolute risk relative to a peer group matched for age and gender), modifiable risk (level of 5-year risk that can be altered by behavior), and absolute risk (probability of disease expression within 5 years). The KYN patient questionnaire was completed online within the 2 weeks preceding the health immersion (baseline) and during

the 2 weeks before the follow-up appointment 7 to 9 months later (endpoint). The biological measures included in the KYN profile were collected in a fasting state on the first day of the immersion and again at the follow-up appointment. Approximately 2 weeks after baseline and endpoint, participants received information on their 5-year disease risk.

Biological Measures—Biological outcome measures included blood pressure (BP), resting pulse, height, weight, body composition, body mass index (BMI), waist circumference, fasting glucose, lipid profile, and high sensitivity C-reactive protein (CRP). BP was measured on the left arm, using a manual sphygmomanometer.^{31,32} Participants were directed to sit quietly for 5 minutes before the first measurement. The radial pulse was taken for 30 seconds after completing the BP, and a second BP and pulse were taken 2 minutes later. Height was measured twice consecutively. Duplicate measurements of pulse, blood pressure, and height were averaged. Body weight was taken on a Tanita Scale (TBF-310GF; Tanita Corporation, Arlington Heights, Illinois) with participants in light indoor clothing and without shoes; participants were asked to fast 12 hours prior to measurements and void before weighing. Body mass index was calculated as weight in kg divided by height in meters squared. Waist circumference was measured in triplicate and averaged according to the method used by Canoy et al.³⁴ All blood samples were collected after participants had fasted for 12 hours. Blood assays were carried out by the Clinical Laboratory Improvement Amendments–certified Duke University Health System core laboratory.

Self-reported Psychosocial Measures

Burns Brief Mood Survey (BMS): The BMS³⁵ includes three five-item subscales for assessment of depression, anxiety, and anger during the previous 1-week period. The BMS has been shown to be valid and reliable.³⁵⁻³⁷ In the present study, Cronbach's α for anger, anxiety, and depression subscales were 0.82, 0.85, and 0.90, respectively.

Relationship Satisfaction Scale (RSAT): The RSAT assesses the level of satisfaction or dissatisfaction felt by the participant in reference to their closest personal relationship; it was selected for the current study as an indicator of social support. This five-item scale demonstrates good convergent and discriminate validity.³⁵ Responses were measured on a Likert scale with anchors of satisfied and dissatisfied. Cronbach's α was 0.97 in the present study.

Perceived Stress Scale (PSS): Participants' nonspecific, self-appraised stress was measured using the four-item PSS. The PSS measures perceived stress over the past month; it has demonstrated reliability and construct validity.^{38,39} Cronbach's α was 0.78 in the present study.

Behavioral Activation

Patient Activation Measure (PAM): The 13-item PAM was used to assess patients' knowledge, skill, and confidence toward self-management of health. Higher scores on this measure have consistently been associated with healthy behavior, health information seeking, and readiness to change in both healthy and chronically ill populations.⁴⁰⁻⁴³ The PAM has been shown to be both valid and reliable.⁴¹ Cronbach's α was 0.77 in the present study.

Readiness to Change (RTC): A six-item RTC questionnaire was designed for this study to assess readiness to change⁴⁴ in the areas of weight reduction, exercise, smoking cessation, diet, stress management, and meditation. The items have independently demonstrated construct validity in previous studies.⁴⁵⁻⁴⁸

Exercise Behavior: Exercise behavior was measured with a brief questionnaire that assessed frequency of aerobic exercise (20 minutes duration), stretching (15 minutes duration), and muscular strengthening exercise (20 minutes duration). Available responses included fewer than once per month, one to two times per week, three to four times per week, and five or more times per week.

Intervention Integrity Check

Wheel of Health Questionnaire (WHQ): The 20-item WHQ, designed for this study, was used to assess whether participants integrated aspects of the intervention into multiple domains of living. Measured dimensions included movement and exercise, nutrition, physical environment, relationships, personal growth and spirituality, mind-body connection, preventive medicine, conventional and CAM treatments, pharmaceuticals/supplements, and mindfulness. Performance and satisfaction in each of the 10 dimensions of the Wheel of Health were measured on a 1-to-10 scale anchored by the descriptors “low” and “high.” All dimensions were summed to create subscores for WHQ performance and WHQ satisfaction.

ANALYSIS

An intention-to-treat analysis was conducted, based upon the 63 participants who were enrolled in the study, attended the 3-day immersion, and completed any assessment at baseline. The last observation was carried forward to replace endpoint values for those participants with missing data. Baseline and endpoint data were excluded (pairwise) in the cases where a participant's baseline questionnaire was incomplete. Analyses were performed using SPSS version 17 (SPSS, Chicago, Illinois). All data except for total cholesterol and HDL cholesterol were non-normally distributed as measured by the Kolmogorov-Smirnov test with Lilliefors Significance Correction and Shapiro-Wilk test ($P < .05$)^{49,50}; therefore, all P -values were determined using the nonparametric Wilcoxon Signed Rank test.⁵¹ Statistical significance was set at the $P < .05$ level.

RESULTS

Participants

Participants ranged in age from 33 to 73 years with a mean age of 59.6 (SD = 8.2). The sample was near equally divided between men and women and consisted predominantly of white individuals who reported relatively high levels of education and income. Thirty-three percent of participants had metabolic syndrome at baseline as defined by the International Diabetes Federation.⁵² Descriptive statistics on the demographic characteristics of this sample are listed in Table 4.

Retention and Adherence

During the support phase, participants completed a median of two of the five available health coaching calls and used an average of 101.0 minutes of total coaching per person (50.5% of available session time). Six of the initial 63 participants were lost to contact or unavailable for the endpoint data collection.

Disease Risk and Disease Risk Biomarkers

Relative Risk: When matched for gender and age with a national sample, the median 5-year relative risk percentile at baseline was well below average at 7% (SD = 20.1%) for coronary heart disease, 16% (SD = 24.1%) for diabetes, and 25% (SD = 28.8%) for stroke.

Modifiable Risk: Five-year median modifiable risk of diabetes shifted ($P = .02$) from 0.37% (SD = 2.55%) at baseline to 0.16% (SD = 2.98%) at endpoint (Figure 2); median stroke risk decreased ($P < .01$) from 0.37% (SD = 2.09%) to 0.23% (SD = 2.32%). Median CHD risk declined from 0.95% (SD = 1.30%) to 0.72% (SD = 1.97%). Although the reduction in median CHD risk is nearly equal in magnitude to that of diabetes and stroke risk, the change in coronary risk was not statistically significant ($P = .17$) due in part to an upward shift in CHD risk variability (interquartile range of 0.23-1.73 at baseline to 0.20-1.87 at endpoint).

Absolute Risk: The change in 5-year absolute risk of diabetes was nonsignificant with a decrease in median risk from 1.28% (SD = 2.87%) at baseline to 1.06% (SD = 3.31%) at endpoint. Median absolute risk of CHD also declined slightly (nonsignificant) from 1.91% (SD = 2.42%) to 1.82% (SD = 2.96%). In contrast, the absolute risk of stroke decreased significantly ($P = .02$) from a median of 1.28% (SD = 2.87%) at baseline to 1.06% (SD = 3.30%) at endpoint.

Biomarkers for disease risk, including BMI ($P = .008$), waist circumference ($P = .003$), and pulse ($P < .001$) improved significantly across the study period (Table 5). Total cholesterol increased significantly during the study ($P = .007$). Systolic blood pressure ($P = .237$), diastolic blood pressure ($P = .095$), HDL cholesterol ($P = .109$), glucose ($P = .300$), and CRP ($P = .068$) did not change.

These data represent intention-to-treat analyses ($n = 63$); however, analyses including only completers ($n = 57$) yielded similar results (data not shown) for all outcomes measured.

Behavioral and Psychosocial: Participants experienced a highly significant increase ($P < .001$) in activation (PAM) during the study period (Table 6), and measures of psychosocial function, including anger, anxiety, depression, and relationship satisfaction, improved significantly ($P < .05$). Importantly, baseline activation (PAM) in completers was not significantly correlated with change in risk of stroke (bivariate $r = 0.09$), diabetes (bivariate $r = 0.20$), and CHD (bivariate $r = -0.12$).

Analyses also demonstrated highly significant ($P < .001$) improvements in the median frequency of aerobic exercise and readiness to change (RTC) for stress reduction and mental focus. Strengthening exercise and exercise stage of change remained unchanged, although all other behavioral measures also improved significantly ($P < .05$) (Table 7). Finally, measures of intervention integrity, including WHQ health performance and satisfaction increased significantly ($P < .001$), as shown in Tables 6 and 7.

DISCUSSION

To our knowledge, this is the first study to evaluate an integrative health model that combines an intensive 3-day health immersion and personalized health plan with follow-up education and physician and coaching support. Integrative medicine builds upon concepts of self-determination theory and emphasizes the individual's role in health.⁵³ For a patient-centered program, one of the key objectives of the current intervention was to help individuals establish health and wellness goals that were based upon their own personal values. A second and equally important objective was to partner with patients in achieving self-stated goals.

Our findings indicate that although cardiac risk remained unchanged, modifiable risk factors for diabetes and stroke decreased significantly across the intervention. While the magnitude of change may appear less than significant from an epidemiological standpoint, it is notable that participants demonstrated improvements in spite of floor effects commonly observed

among relatively healthy populations. In addition, multiple psychosocial and behavioral measures improved and participants became more activated towards self-management of health. Significant improvements in patient activation indicate that participants became more confident in making and maintaining healthy lifestyle changes. Importantly, baseline activation did not mediate the changes in modifiable risk. Rather, the intervention itself appears to have activated patients to change lifestyle behavior and reduce risk.

The mean BMI of participants at baseline was near the normal range at 26.8, so many patients sought to maintain an already healthy weight rather than include weight loss among their health goals. Maintenance of a healthy weight represents a clinically relevant outcome in and of itself since body mass tends to increase through the sixth decade of life and such increases are related to high absolute risk of disease and mortality.⁵⁴ Participants lost 0.9 kg (approximately 2 lbs; mean = 80.5, SD = 21.3 to mean = 79.6 SD = 21; $P = .01$) across the intervention; however, even moderate weight loss has been associated with improvements in risk factors for cardiovascular disease and diabetes.⁵⁵⁻⁵⁷ Furthermore, each kilogram of weight loss is on average associated with a decrease in incidence of diabetes by 16%⁵⁸ and a reduction in systolic blood pressure of 1.0 to 2.4 mm Hg.⁵⁹

While short-term improvements in KYN disease risk measures may appear modest, the KYN model did not account for the potential long-term benefits that can result from reduced anger, anxiety, and depression and increased social support. Such psychosocial and quality of life measures have shown utility in predicting future health status and mortality.^{60,61} High levels of anger, anxiety, and depression have been associated with risk of cardiovascular disease⁶² and stroke.⁶³ Hence, the reduction in anger, anxiety, and depression and improvement in WHQ satisfaction and performance may represent long-term benefits not captured by the KYN disease risk model.

Changes Accounting for Reduction in Disease Risk

A significant drop in mean BMI and waist circumference and an increase in aerobic exercise frequency (Tables 5 and 6) explain the decrease in modifiable diabetes risk, as calculated by the KYN disease assessment tool (Table 3). Improvements in BMI and exercise frequency also accounted for the drop in modifiable risk of stroke after the intervention. In contrast, the CHD risk-reducing effects of decreased BMI and increased exercise level were counterbalanced by increases in total cholesterol; these counterbalanced effects explain the lack of improvement in modifiable CHD risk.

The increase in cholesterol observed in the study was interesting since total cholesterol does not increase appreciably with age during the fifth and sixth decades of life.⁶⁴ It is not clear which factors contributed to the mean increase in cholesterol; however, a portion of the change can be attributed to three participants whose total cholesterol increased by more than 79 mg/dL each. One of these was diagnosed with cancer and experienced a notable increase in body mass, the second participant increased weight by 1.2 kg (2.6 lbs) and discontinued a lipid lowering medication, and the third participant gained 0.9 kg (2 lbs) but otherwise reported no notable changes.

Resting pulse was not part of the disease risk algorithm; nonetheless, it decreased significantly between baseline and endpoint, probably related to improved exercise level reflecting better fitness and cardiovascular conditioning.

Patient activation at baseline did not predict the variability in disease risk reduction. This result is consistent with the intervention's personalized approach to lifestyle change. The health immersion and follow-up coaching were tailored to the participant's baseline level of motivation and engagement. Clinicians and coaches worked within this context to enhance

patient engagement and accomplish patient-stated goals. These results are encouraging from the standpoint that participants were able to achieve disease risk reduction regardless of baseline level of activation. The increases in PAM seen in the present study can be benchmarked against a previous intervention that also used tailored telephonic coaching as a disease management strategy. Both studies demonstrated significant increases in mean PAM scores; however, the current study showed an increase in score of 8.4 (68.3 at baseline to 76.7 at endpoint) compared to an increase of 4.6 (64.3 at baseline to 68.9 at endpoint) in the benchmark trial.⁶⁵

The moderate utilization of telephonic coaching (50.5% mean utilization of available coaching) is of interest. It contrasts with two of our other studies that used the same approach and even some of the same coaches; in these two RCT studies, telephonic coaching adherence was 74%²⁶ and 93%.⁶⁶ Possible explanations for the utilization differences in the present study include the following: (1) The study intervention offered “optional” IH coaching whereas coaching was much more strongly encouraged in the other two intervention studies; (2) distinct demographics in the present study (ie, older population with fewer women, minorities, and persons of low or moderate socioeconomic status) compared to the previous trials; and (3) according to the treatment team and lead physician, the present sample was composed of “high achievement-oriented individuals, many of whom wanted a personalized health plan and then acted on it independently.” Hence, it appears that latent factors such as motivation, personality (eg, Type A), and life experiences (many executives) contributed to the lower level of coaching utilization.

The improvements in disease risk biomarkers seen here are somewhat less extensive than those reported from intensive multimodal lifestyle interventions such as those conducted by Daubenmier et al²³ and Ornish et al.²⁴ There are, however, substantive differences between the present intervention and previous lifestyle trials. Namely, the current intervention was significantly shorter in duration and intensity. In addition, the current trial was conducted among a sample with lower initial disease risk. Given the rapidly growing cost concerns in health care, it is reasonable to evaluate outcomes in the context of program duration and intensity. Within this context, the results of the current study are notable since they were achieved with a 3-day intervention plus fewer than 4 hours of telephonic follow-up. In contrast, the Lifestyle Heart Trial conducted by Ornish et al²⁴ employed a 7-day retreat with 4-hour twice-weekly group support meetings lasting for 1 year.

The lifestyle changes in the present study are comparable to those of a previous controlled trial that tested the effects of an integrative intervention with health coaching²⁶ on cardiovascular risk. The current study was conducted in a healthier population, utilized less coaching, and capitalized on the added benefit of a 3-day health immersion. Both studies, however, demonstrated similar baseline-to-endpoint improvements in readiness to lose weight and increase physical activity, frequency of exercise, and reduction in BMI.

Future study of integrative health models of care would benefit from RCT designs with larger samples. There is also a need to further explore the mechanisms of behavior change and the dose effect of these interventions. Finally, from a cost-effectiveness perspective, it may be useful to clarify which aspects of the program are most effective for which participants.

The principle limitations of this study were low sample diversity, moderate sample size, and the observational study design. The intervention demonstrated effectiveness among a sample of relatively healthy older adults who were predominantly white and reported income and education well above the median. It is yet to be seen if these findings will generalize to other populations. A self-selection bias may have also influenced the results; participants who

elected to take part in this study may have had stronger motivation to change health behaviors. However, selection bias is not evident based on initial patient activation scores, which are in fact slightly below the normative mean of 69 observed among employed adults.⁴⁰ Also, the observational study design, though appropriate for exploratory purposes, does not account for the influence of expectancy or the natural course of health. Finally, analysis of mean changes in heterogeneous groups presents a challenge since large sample variability can mask important shifts in subsets of individuals. This warrants further study utilizing methods such as structural equation modeling and path analysis to assess individual differences and determine patterns of change.

CONCLUSION

Health behavior changes were accompanied by an improvement in patient activation and psychosocial measures (anger, anxiety, depression, relationship satisfaction) and a decrease in modifiable risk of stroke and diabetes. This suggests that an integrative health model can help patients become more engaged in self-management of health and support them in making and maintaining healthy lifestyle changes.

Acknowledgments

The authors would like to thank Shin-Yiing Yeung, BS; Tracey Hawkins, BA; Lesley Chapman, BS; and Marius Kluenger, BS; for their assistance in data collection; Julie D. Pruitt, MS, RD; Bill Kraus, MD; Connie Bales, PhD; and ActivHealth for the use of their online health education materials; Timothy Smith, PhD; Mark Ruby; Guizhou Hu, MD, PhD; and Biosignia for technical assistance and use of the Know Your Number assessment; Linda Smith, PA, for intervention program oversight; Janna Fikkan, PhD; and Mark Dreusicke, BS; for manuscript editing; the Cliffs Communities for financial and recruitment support; Holbrook Raynal, MD, DHA, and the Preserve Health team for providing facilities and support for endpoint data collection; the Ripple Foundation for infrastructure funding of the Duke Integrative Medicine (IM) research team; and the Duke IM clinicians who supported this program: Sam Moon, MD, MPH; Shelley Wroth, MD; Evangeline Lausier, MD; Louise Goldstein, RN; Jeff Brantley, MD; Beth Reardon, MS, RD; Janet Shaffer, LAc; Jennifer Davis, MA, LPC; Jeanne Van Gemert, MA, LPC; Jessica P. Wakefield, MA, LPC; Julie Kosey, MS, CPCC; Kerry Little, MA, CPCC; JoAnna Stange, BA, CPCC; Faith Seay, RN; Mary Peterson, MA; and Noah St John, MS. This study was funded in part by The Cliffs Communities, Inc, and in part by the Ripple Foundation. The paper was supported by grant K99 AT004945 from the National Institutes of Health (NIH) National Center for Complementary and Alternative Medicine to J. M. Greeson, and its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIH.


REFERENCES

1. Lloyd-Jones D, Adams R, Carnethon M, et al. Heart disease and stroke statistics—2009 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation*. 2009; 119(3):e21–e181. [PubMed: 19075105]
2. Van Dam RM, Rimm EB, Willett WC, Stampfer MJ, Hu FB. Dietary patterns and risk for type 2 diabetes mellitus in U.S. men. *Ann Intern Med*. 2002; 136(3):201–209. [PubMed: 11827496]
3. Eyre H, Kahn R, Robertson RM, et al. American Cancer Society; American Diabetes Association; American Heart Association. Preventing cancer, cardiovascular disease, and diabetes: a common agenda for the American Cancer Society, the American Diabetes Association, and the American Heart Association. *Stroke*. 2004; 35(8):1999–2010. [PubMed: 15272139]
4. Kromhout D, Menotti A, Kesteloot H, Sans S. Prevention of coronary heart disease by diet and lifestyle: evidence from prospective cross-cultural, cohort, and intervention studies. *Circulation*. 2002; 105(7):893–898. [PubMed: 11854133]
5. American Heart Association Nutrition Committee. Lichtenstein AH, Appel LJ, Brands M, et al. Diet and lifestyle recommendations revision 2006: a scientific statement from the American Heart Association Nutrition Committee. *Circulation*. 2006; 114(1):82–96. [PubMed: 16785338]
6. Suiitor CW, Meyers LD, Institute of Medicine. Dietary Reference Intakes Research Synthesis Workshop summary. 2007

7. Bantle JP, Wylie-Rosett J, Albright AL, et al. Nutrition recommendations and interventions for diabetes—2006: a position statement of the American Diabetes Association. *Diabetes Care*. 2006; 29(9):2140–2157. [PubMed: 16936169]
8. Bunker SJ, Colquhoun DM, Esler MD, et al. “Stress” and coronary heart disease: psychosocial risk factors. *Med J Aust*. 2003; 178(6):272–276. [PubMed: 12633484]
9. Pleis JR, Lucas JW. Summary health statistics for U.S. adults: National Health Interview Survey, 2007. *Vital Health Stat* 10. May.2009 (240):1–159.
10. Centers for Disease Control and Prevention (CDC). Trends in intake of energy and macronutrients —United States, 1971–2000. *MMWR Morb Mortal Wkly Rep*. 2004; 53(4):80–82. [PubMed: 14762332]
11. Uchino BN. Social support and health: a review of physiological processes potentially underlying links to disease outcomes. *J Behav Med*. 2006; 29(4):377–387. [PubMed: 16758315]
12. Marks R, Allegrante JP, Lorig K. A review and synthesis of research evidence for self-efficacy-enhancing interventions for reducing chronic disability: implications for health education practice (part II). *Health Promot Pract*. 2005; 6(2):148–156. [PubMed: 15855284]
13. Dyer J. How does spirituality affect physical health? A conceptual review. *Holist Nurs Pract*. 2007; 21(6):324–328. [PubMed: 17978636]
14. Carmody J, Reed G, Kristeller J, Merriam P. Mindfulness, spirituality, and health-related symptoms. *J Psychosom Res*. 2008; 64(4):393–403. [PubMed: 18374738]
15. Serra-Majem L, Roman B, Estruch R. Scientific evidence of interventions using the Mediterranean diet: a systematic review. *Nutr Rev*. 2006; 64(2 Pt 2):S27–S47. [PubMed: 16532897]
16. Penedo FJ, Dahn JR. Exercise and well-being: a review of mental and physical health benefits associated with physical activity. *Curr Opin Psychiatry*. 2005; 18(2):189–193. [PubMed: 16639173]
17. Innes KE, Bourguignon C, Taylor AG. Risk indices associated with the insulin resistance syndrome, cardiovascular disease, and possible protection with yoga: a systematic review. *J Am Board Fam Med*. 2005; 18(6):491–519.
18. Yang K. A review of yoga programs for four leading risk factors of chronic diseases. *Evid Based Complement Alternat Med*. 2007; 4(4):487–491. [PubMed: 18227916]
19. Greeson JM. Mindfulness research update: 2008. *Complement Health Pract Rev*. 2009; 14(1):10–18. [PubMed: 20047019]
20. Anderson JW, Liu C, Kryscio RJ. Blood pressure response to transcendental meditation: a meta-analysis. *Am J Hypertens*. 2008; 21(3):310–316. [PubMed: 18311126]
21. Paul-Labrador M, Polk D, Dwyer JH, et al. Effects of a randomized controlled trial of transcendental meditation on components of the metabolic syndrome in subjects with coronary heart disease. *Arch Intern Med*. 2006; 166(11):1218–1224. [PubMed: 16772250]
22. Wolever RQ, Caldwell KL, Wakefield JP, et al. Integrative Health Coaching: An organizational case study. *Explore*. 2011; 7(1):30–36. [PubMed: 21194670]
23. Daubenmier JJ, Weidner G, Sumner MD, et al. The contribution of changes in diet, exercise, and stress management to changes in coronary risk in women and men in the multisite cardiac lifestyle intervention program. *Ann Behav Med*. 2007; 33(1):57–68. [PubMed: 17291171]
24. Ornish D, Brown SE, Scherwitz LW, et al. Can lifestyle changes reverse coronary heart disease? The Lifestyle Heart Trial. *Lancet*. 1990; 336(8708):129–133. [PubMed: 1973470]
25. The Center for Integrative Medicine at Duke University. Liebowitz, R.; Smith, L.; Gaudet, T. *The Duke Encyclopedia of New Medicine: Conventional and Alternative Medicine for All Ages*. Rodale; London, England; New York, NY: 2006. p. 24
26. Edelman D, Oddone EZ, Liebowitz RS, et al. A multidimensional integrative medicine intervention to improve cardiovascular risk. *J Gen Intern Med*. 2006; 21(7):728–734. [PubMed: 16808774]
27. Cooperrider, D.; Whitney, D. *Appreciative Inquiry: A Positive Revolution in Change*. Berrett-Koehler Publishers; San Francisco, CA: 2005.
28. Miller, WR.; Rollnick, S. 2nd ed.. Guilford Press; New York, New York: 2002. *Motivational Interviewing*.

29. Hu G, Root MM. Building prediction models for coronary heart disease by synthesizing multiple longitudinal research findings. *Eur J Cardiovasc Prev Rehabil*. 2005; 12(5):459–464. [PubMed: 16210932]
30. Samsa G, Hu G, Root M. Combining information from multiple data sources to create multivariable risk models: illustration and preliminary assessment of a new method. *J Biomed Biotechnol*. 2005; 2005(2):113–123. [PubMed: 16046816]
31. Beevers G, Lip GYH, O'Brien E. ABC of hypertension. Blood pressure measurement. Part I—sphygmomanometry: factors common to all techniques. *Br Med J*. 2001; 322(7292):981–985. [PubMed: 11312235]
32. Pickering TG, Hall JE, Appel LJ, et al. Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. Recommendations for blood pressure measurement in humans and experimental animals. Part 1: Blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Hypertension*. 2005; 45(1):142–161. [PubMed: 15611362]
33. Jebb SA, Cole TJ, Doman D, Murgatroyd PR, Prentice AM. Evaluation of the novel Tanita body-fat analyser to measure body composition by comparison with a four-compartment model. *Br J Nutr*. 2000; 83(02):115–122. [PubMed: 10743490]
34. Canoy D, Boekholdt SM, Wareham N, et al. Body fat distribution and risk of coronary heart disease in men and women in the European Prospective Investigation Into Cancer and Nutrition in Norfolk cohort: a population-based prospective study. *Circulation*. 2007; 116(25):2933–2943. [PubMed: 18071080]
35. Burns, DD. *Therapist's Toolkit: Comprehensive Assessment and Treatment Tools for the Mental Health Professional*. Author; Gladwyne, PA: 1995.
36. Sekirnjak, GC.; Beal, D. The concurrent validity of the Burns Depression Checklist.. Annual Meeting of the Association for the Advancement of Behavior Therapy.; Toronto, Canada. November 1999;
37. Sekirnjak, GC.; Beal, D. The Concurrent Validity of the Burns Anxiety Checklist.. Annual Meeting of the Association for the Advancement of Behavior Therapy.; Toronto, Canada. November 1999;
38. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav*. 1983; 24(4):385–396. [PubMed: 6668417]
39. Pbert L, Doerfler LA, DeCosimo D. An evaluation of the perceived stress scale in two clinical populations. *J Psychopathol Behav Assess*. 1992; 14(4):363–375.
40. Fowles JB, Terry P, Xi M, Hibbard J, Bloom CT, Harvey L. Measuring self-management of patients' and employees' health: further validation of the Patient Activation Measure (PAM) based on its relation to employee characteristics. *Patient Educ Couns*. 2009; 77(1):116–122. [PubMed: 19356881]
41. Hibbard JH, Mahoney ER, Stockard J, Tusler M. Development and testing of a short form of the patient activation measure. *Health Serv Res*. 2005; 40(6 Pt 1):1918–1930. [PubMed: 16336556]
42. Hibbard JH, Mahoney ER, Stock R, Tusler M. Do increases in patient activation result in improved self-management behaviors? *Health Serv Res*. 2007; 42(4):1443–1463. [PubMed: 17610432]
43. Mosen DM, Schmittiel J, Hibbard J, Sobel D, Remmers C, Bellows J. Is patient activation associated with outcomes of care for adults with chronic conditions? *J Ambul Care Manage*. 2007; 30(1):21–29. [PubMed: 17170635]
44. Prochaska JO, Velicer WF. The transtheoretical model of health behavior change. *Am J Health Promot*. 1997; 12(1):38–48. [PubMed: 10170434]
45. Marshall SJ, Biddle SJ. The transtheoretical model of behavior change: a meta-analysis of applications to physical activity and exercise. *Ann Behav Med*. 2001; 23(4):229–246. [PubMed: 11761340]
46. Greene GW, Rossi SR, Rossi JS, Velicer WF, Fava JL, Prochaska JO. Dietary applications of the stages of change model. *J Am Diet Assoc*. 1999; 99(6):673–678. [PubMed: 10361528]

47. DiClemente CC, Prochaska JO, Fairhurst SK, Velicer WF, Velasquez MM, Rossi JS. The process of smoking cessation: an analysis of precontemplation, contemplation, and preparation stages of change. *J Consult Clin Psychol*. 1991; 59(2):295–304. [PubMed: 2030191]
48. Nigg CR, Burbank PM, Padula C, et al. Stages of change across ten health risk behaviors for older adults. *Gerontologist*. 1999; 39(4):473–482. [PubMed: 10495586]
49. Shapiro SS, Wilk MB. An analysis of variance test for normality (complete samples). *Biometrika*. 1965; 52(3-4):591–611.
50. Lilliefors HW. On the Kolmogorov-Smirnov test for normality with mean and variance unknown. *J Am Stat Assoc*. 1967:399–402.
51. Wilcoxon F. Individual comparisons by ranking methods. *Biometrics Bull*. 1945; 1(6):80–83.
52. Alberti KG, Zimmet P, Shaw J, IDF Epidemiology Task Force Consensus Group. The metabolic syndrome—a new worldwide definition. *Lancet*. 2005; 366(9491):1059–1062. [PubMed: 16182882]
53. Ryan RM, Deci EL. Self-determination theory and the facilitation of intrinsic motivation, social development, and well-being. *Am Psychol*. 2000; 55(1):68–78. [PubMed: 11392867]
54. Nooyens AC, Visscher TLS, Verschuren WM, et al. Age, period and cohort effects on body weight and body mass index in adults: The Doetinchem Cohort Study. *Public Health Nutr*. 2009; 12(06): 862–870. [PubMed: 18652715]
55. Knowler WC, Barrett-Connor E, Fowler SE, et al. Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med*. 2002; 346(6):393–403. [PubMed: 11832527]
56. Stevens VJ, Obarzanek E, Cook NR, et al. Trials for the Hypertension Prevention Research Group. Long-term weight loss and changes in blood pressure: results of the Trials of Hypertension Prevention, phase II. *Ann Intern Med*. 2001; 134(1):1–11. [PubMed: 11187414]
57. Tuomilehto J, Lindstrom J, Eriksson JG, et al. Finnish Diabetes Prevention Study Group. Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med*. 2001; 344(18):1343–1350. [PubMed: 11333990]
58. Hamman RF, Wing RR, Edelstein SL, et al. Effect of weight loss with lifestyle intervention on risk of diabetes. *Diabetes Care*. 2006; 29(9):2102–2107. [PubMed: 16936160]
59. Neter JE, Stam BE, Kok FJ, Grobbee DE, Geleijnse JM. Influence of weight reduction on blood pressure: a meta-analysis of randomized controlled trials. *Hypertension*. 2003; 42(5):878–884. [PubMed: 12975389]
60. Wolinsky FD, Johnson RJ. Perceived health status and mortality among older men and women. *J Gerontol*. 1992; 47(6):S304–S312. [PubMed: 1430868]
61. Spertus JA, Jones P, McDonnell M, Fan V, Fihn SD. Health status predicts long-term outcome in outpatients with coronary disease. *Circulation*. 2002; 106(1):43–49. [PubMed: 12093768]
62. Suls J, Bunde J. Anger, anxiety, and depression as risk factors for cardiovascular disease: the problems and implications of overlapping affective dispositions. *Psychol Bull*. 2005; 131(2):206–300.
63. Rutledge T, Hogan BE. A quantitative review of prospective evidence linking psychological factors with hypertension development. *Psychosom Med*. 2002; 64(5):758–766. [PubMed: 12271106]
64. Abbott RD, Sharp DS, Burchfiel CM, et al. Cross-sectional and longitudinal changes in total and high-density-lipoprotein cholesterol levels over a 20-year period in elderly men: the Honolulu Heart Program. *Ann Epidemiol*. 1997; 7(6):417–424. [PubMed: 9279451]
65. Hibbard JH, Greene J, Tusler M. Improving the outcomes of disease management by tailoring care to the patient's level of activation. *Am J Manag Care*. 2009; 15(6):353–360. [PubMed: 19514801]
66. Wolever RQ, Dreusicke MH, Fikkan JL, et al. Integrative Health Coaching for patients with type 2 diabetes: a randomized clinical trial. *Diabetes Educ*. 2010; 36(4):629–639. [PubMed: 20534872]



*Complementary and alternative medicine treatments

FIGURE 1. Duke Integrative Medicine Wheel of Health
Used with permission from Duke Integrative Medicine for the Wheel of Health.




FIGURE 2. Median 5-year Modifiable Disease Risk

The median modifiable risk of diabetes decreased (* $P=.02$) from 0.37% SD 2.55% to 0.16% SD 2.98%; risk of stroke decreased significantly (** $P<.01$) from 0.37% SD 2.09% to 0.23% SD 2.32%; and risk of coronary heart disease declined (nonsignificant; $P=.17$) from 0.95% SD 1.30% to 0.72% SD 1.97%.

TABLE 1**Duke Integrative Medicine Three-day Health Immersion Program**

<ol style="list-style-type: none"> 1. Introduction to Integrative Medicine, information on the Three-day Health Immersion program, description of personalized health planning, and explanation of the Duke Integrative Medicine Wheel of Health 2. Individual integrative physician consultations <ul style="list-style-type: none"> • New patient consultation on day 1 (80 min) • Follow-up consultation on day 3 (30 min) 3. Health coaching <ul style="list-style-type: none"> • Three group coaching sessions (50 min each) or one individual coaching session via phone (90 min) 4. Individual fitness assessment (60 min) 5. Individual manual therapies <ul style="list-style-type: none"> • Acupuncture or acupressure (60 min) • Massage (60 min) 6. Group movement classes <ul style="list-style-type: none"> • Yoga (60 min) • Qigong (60 min) 7. Nutrition consultation and education <ul style="list-style-type: none"> • Individual integrative nutrition consult with registered dietician (60 min) • Nutrition class: In Search of the Optimal Diet (60 min) 8. Healthful cooking class (60 min) 9. Individual mind/body consultation (60 min) 10. Group mindfulness classes <ul style="list-style-type: none"> • Introduction to mindfulness-based meditation (60 min) • Mindful-eating lunch (60 min) 11. Program conclusion and final reflection (group setting)

All subjects participated in the above health-related activities during a 3-day period between January and March 2008.

TABLE 2

Group Education Sessions Provided During the Support Phase

1. A Healing Journey Through Cancer: Integrative Medicine Approaches for Prevention and Treatment
2. Maximize Back, Joint, and Muscular Health
3. A Woman's Midlife Health Journey: Embracing Perimenopause and Menopause
4. When Working Out Hurts: Alternative Treatments for Pain Management and Relief for Sports and Exercise-related Injuries
5. Integrative Medicine Strategies for Optimal Living: Approaches to the Prevention and Treatment of Cardiovascular Disease
6. Give Yourself Health: The Power of Your Mind/Body Connection
7. In Search of the Optimal Diet
8. Emotional Wellness: How Emotions Affect Your Health and Well-being

TABLE 3**Know Your Number Disease Risk Components***

Risk Factor	CHD	Stroke	Diabetes
Age			
Gender			
Ethnicity			
Diagnosed with CHD			
Previous stroke			
Diagnosed with diabetes			
Diagnosed with LVH			
Family history of CHD			
Daily aspirin use			
Current HRT therapy			
BMI			
Systolic blood pressure			
Current smoker			
Past smoker			
Exercise level			
Total cholesterol			
HDL			
C-reactive protein			
Diagnosed with other CVD			
Diagnosed with atrial fibrillation			
Family history of stroke			
Current use of hypertension medication			
Previous diagnosis of gestational diabetes			
Years since gestational diabetes			
Family history of diabetes			
Waist measurement			
Fasting blood glucose			

Abbreviations: CHD, coronary heart disease; LVH, left-ventricular hypertrophy; HRT, hormone replacement therapy; BMI, body mass index; HDL, high-density lipoprotein; CVD, cardiovascular disease.

* Diabetes indicates type 2 diabetes mellitus; darkened blocks reflect variables used to calculate specific disease risks. Table content provided by Biosignia.

TABLE 4

Participant Baseline Demographics (n = 63)

Highest Level of Education	
Grade school	1 (1.6%)
High school	3 (6.3%)
Undergraduate	23 (36.5%)
Graduate	22 (34.9%)
Postgraduate	14 (22.2%)
Gender	
Male	33 (52.4%)
Female	30 (47.6%)
Ethnicity	
White non-Latino	62 (98.4%)
Other	1 (1.6%)
Age, y	
<40	1 (1.6%)
40-49	8 (12.7%)
50-59	15 (23.8%)
>60	39 (61.9%)
Marital status	
Single	5 (7.9%)
Married	55 (87.3%)
Widowed	3 (4.8%)
Annual household income	
\$20 001-\$50 000	2 (3.5%)
\$50 001-\$100 000	7 (12.3%)
>\$100 000	48 (84.2%)
Work status	
Employed full-time	18 (28.6%)
Employed part-time	3 (4.8%)
Unemployed	1 (1.6%)
Retired	40 (63.5%)
Other	1 (1.6%)
Health status	
Metabolic syndrome *	21 (33.3%)

* International Diabetes Federation criteria for metabolic syndrome.

TABLE 5

Changes in Biomarkers of Disease Risk With Integrative Program*

	Baseline			Endpoint			P value [†]
	Mean	Median	SD	Mean	Median	SD	
BMI (kg/m ²)	26.8	26.7	5.5	26.5	25.8	25.8	.008
Waist circumference (cm)	91.6	90.0	15.3	90.2	89.5	89.5	.003
Pulse (BPM)	64.8	64.0	9.9	60.7	60.0	60.0	<.001
Systolic BP (mm Hg)	121.0	119.5	13.7	119.8	116.0	116.0	.237
Diastolic BP (mm Hg)	76.8	78.0	7.4	75.6	75.0	75.0	.095
Total cholesterol (mg/dL)	198.8	195.0	40.1	212.4	205.0	205.0	.007
HDL cholesterol (mg/dL)	65.4	66.0	21.0	63.1	64.0	64.0	.109
Glucose (mg/dL)	95.5	91.0	17.1	96.3	95.0	95.0	.300
CRP (mg/L)	2.3	1.4	3.8	2.2	1.2	1.2	.068

Abbreviations: BMI, body mass index; BPM, beats per minute; BP, blood pressure; HDL, high-density lipoprotein; CRP, C-reactive protein (high sensitivity).

* Intention-to-treat sample (n = 63).

[†] P-value was calculated by the Wilcoxon Signed Rank Test.

TABLE 6

Changes in Psychosocial Measures With Integrative Program*

	Baseline			Endpoint			P value [†]
	Mean	Median	SD	Mean	Median	SD	
Anger BMS (0-20)	2.6	2.0	2.6	1.9	1.0	2.6	.014
Anxiety BMS (0-20)	3.3	3.0	3.0	2.4	1.0	2.9	.004
Depression BMS (0-20)	2.0	1.0	2.8	1.4	0.0	2.5	.021
Perceived stress PSS (0-16)	3.8	3.0	2.8	4.3	4.0	2.8	.311
Relationship dissatisfaction RSAT (0-30)	21.7	25.0	8.1	24.4	26.0	6.8	.004
Patient activation PAM (0-100)	68.3	68.0	16.3	76.7	78.0	16.1	<.001
Health satisfaction WHQ (10-100)	68.6	69.0	13.1	75.5	76.0	13.3	<.001

Abbreviations: BMS, Brief Mood Survey; PSS, Perceived Stress Scale; RSAT, Relationship Satisfaction Test; PAM, Patient Activation Measure; WHQ, Wheel of Health Questionnaire.

* Intention-to-treat sample (n = 63).

[†] P value was calculated by the Wilcoxon Signed Rank Test.

TABLE 7

Changes in Health Behavior With Integrative Program*

	Baseline			Endpoint			P value [†]
	Mean	Median	SD	Mean	Median	SD	
Stretching frequency (1-4)	1.9	2.0	0.9	2.2	2.0	1.0	.006
Aerobic exercise frequency (1-4) [‡]	2.3	2.0	1.0	2.6	3.0	0.9	<.001
Strengthening exercise frequency (1-4)	1.8	2.0	0.8	1.9	2.0	0.8	.252
State of Change (RTC): Weight reduction (0-4)	1.8	1.0	1.4	2.4	3.0	1.4	.009
State of Change (RTC): Exercise RTC (0-4)	2.5	3.0	1.3	2.7	3.0	1.3	.122
State of Change (RTC): Healthier eating (0-4)	2.7	3.0	1.2	3.0	3.0	1.0	.008
State of Change (RTC): Stress reduction (0-5)	2.7	3.0	1.8	3.4	4.0	1.7	<.001
State of Change (RTC): Mental focus (0-4)	1.7	1.0	1.3	2.4	3.0	1.4	<.001
Health Performance WHQ (10-100)	67.6	68.5	11.7	74.7	77.0	12.7	<.001

Abbreviations: RTC, readiness to change; WHQ, Wheel of Health Questionnaire.

* Intention-to-treat sample (n = 63).

[†] P-value was calculated by the Wilcoxon Signed Rank Test.

[‡] Exercise (>20 min) frequency per week: 1 = fewer than once; 2 = 1-2 times; 3 = 3-4 times; 4 = 5 or more times.